MATH 10005

MULTIPLICATION OF RADICALS

Definition:

• *n*-th root of *a*: The *n*-th root of *a*, denoted $\sqrt[n]{a}$, is a number whose *n*-th power equals *a*. In other words,

$$\sqrt[n]{a=b}$$
 means $b^n=a$.

The number n is called the **index**.

Important Properties:

• Product rule for radicals: If $\sqrt[n]{a}$ and $\sqrt[n]{b}$ are real numbers and n is a positive integer,

$$\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{ab}.$$

In other words, the product of radicals is the radical of the product.

- FOIL: <u>First Outer Inner Last</u>. This is one method for multiplying factors which have two terms.
- Distributive Property: Recall that

$$a(b+c) = ab + ac$$
 and $a(b-c) = ab - ac$

• Special Formulas: Note that you do not need to memorize these formulas. They arise by using FOIL.

$$\boxed{(x+y)^2 = x^2 + 2xy + y^2}$$
$$\boxed{(x-y)^2 = x^2 - 2xy + y^2}$$
$$\boxed{(x-y)(x+y) = x^2 - y^2}$$

Simplifying radicals: A radical is in simplest form when the following conditions are satisfied.

- The quantity under the radical has no factor raised to a power greater than or equal to the index.
- There is no fraction under the radical.
- There is no radical in the denominator.
- There is no common factor, other than 1, between the exponents on factors under the radical and the index.

Common Mistakes to Avoid:

- Do not drop the index when working with cubes and other *n*-th roots.
- Do NOT distribute factors on the outside of the radical inside the radical. For example,

$$3x\sqrt{2x} \neq \sqrt{6x^2}.$$

• Remember $(a+b)^2 \neq a^2 + b^2$.

PROBLEMS

Multiply each product. Write all answers in simplest form.

4. $(\sqrt{8}-3)^2$ 1. $3\sqrt{2}(5\sqrt{3}-4)$ $(\sqrt{8}-3)^2$ $3\sqrt{2}(5\sqrt{3}-4)$ $(\sqrt{8}-3)(\sqrt{8}-3)$ $15\sqrt{6} - 12\sqrt{2}$ $\sqrt{64} - 3\sqrt{8} - 3\sqrt{8} + 9$ $8 - 3\sqrt{8} - 3\sqrt{8} + 9$ $17 - 6\sqrt{8}$ 2. $5x\sqrt[3]{9}(2\sqrt[3]{3}-6\sqrt[3]{9})$ $17 - 6\sqrt{4 \cdot 2}$ $17 - 6\sqrt{4}\sqrt{2}$ $5x\sqrt[3]{9}(2\sqrt[3]{3}-6\sqrt[3]{9})$ $17 - 6 \cdot 2\sqrt{2}$ $10x\sqrt[3]{27} - 30x\sqrt[3]{81}$ $17 - 12\sqrt{2}$ $10x \cdot 3 - 30x\sqrt[3]{27 \cdot 3}$ $30x - 30x\sqrt[3]{27}\sqrt[3]{3}$ $30x - 90x\sqrt[3]{3}$ 5. $(2\sqrt{3}+\sqrt{6})^2$ $(2\sqrt{3}+\sqrt{6})^2$ $(2\sqrt{3}+\sqrt{6})(2\sqrt{3}+\sqrt{6})$ 3. $(3\sqrt{5}+2)(2\sqrt{5}-7)$ $4\sqrt{9} + 2\sqrt{18} + 2\sqrt{18} + \sqrt{36}$ $4 \cdot 3 + 4\sqrt{18} + 6$ $(3\sqrt{5}+2)(2\sqrt{5}-7)$ $12 + 4\sqrt{9}\sqrt{2} + 6$ $6\sqrt{25} - 21\sqrt{5} + 4\sqrt{5} - 14$ $18 + 4 \cdot 3\sqrt{2}$ $6 \cdot 5 - 21\sqrt{5} + 4\sqrt{5} - 14$ $18 + 12\sqrt{2}$ $30 - 21\sqrt{5} + 4\sqrt{5} - 14$ $16 - 17\sqrt{5}$ 6. $(3\sqrt{5}-4)(3\sqrt{5}+4)$ $(3\sqrt{5}-4)(3\sqrt{5}+4)$ $9\sqrt{25} + 12\sqrt{5} - 12\sqrt{5} - 16$ $9 \cdot 5 - 16$

$$45 - 16$$

7. $(\sqrt{7}+2)(\sqrt{2}-4)$

$$\frac{\left(\sqrt{7}+2\right)\left(\sqrt{2}-4\right)}{\sqrt{14}-4\sqrt{7}+2\sqrt{2}-8}$$

8. $(2\sqrt{8} - \sqrt{3})(2\sqrt{48} - \sqrt{2})$

$$\begin{array}{c} \left(2\sqrt{8}-\sqrt{3}\right)\left(2\sqrt{48}-\sqrt{2}\right)\\ \left(2\sqrt{4\cdot 2}-\sqrt{3}\right)\left(2\sqrt{16\cdot 3}-\sqrt{2}\right)\\ \left(2\cdot 2\sqrt{2}-\sqrt{3}\right)\left(2\cdot 4\sqrt{3}-\sqrt{2}\right)\\ \left(4\sqrt{2}-\sqrt{3}\right)\left(8\sqrt{3}-\sqrt{2}\right)\\ 32\sqrt{6}-4\sqrt{4}-8\sqrt{9}+\sqrt{6}\\ 32\sqrt{6}-4\cdot 2-8\cdot 3+\sqrt{6}\\ 32\sqrt{6}-8-24+\sqrt{6}\\ \hline -32+33\sqrt{6} \end{array} \right)$$

NOTE: You could also have multiplied using FOIL first and then simplified your answer.