
Topic 1: Arguments

An **argument** is a set of statements in which one of the statements is called the conclusion and the rest comprise the hypotheses.

A **valid argument** is an argument in which the conclusion must be true whenever the hypothesis is true.

$$\begin{array}{l} \text{If it is snowing, then it is cold.} \\ \text{It is snowing.} \\ \hline \text{Therefore, it is cold.} \end{array}$$

Let p = it is snowing and let q = it is cold. Then symbolically, the above argument can be represented as:

$$\begin{array}{l} p \rightarrow q \\ p \\ \hline q \end{array}$$

We can determine the validity of an argument by considering its corresponding truth table.

p	q	$p \rightarrow q$	$(p \rightarrow q) \wedge p$	$[(p \rightarrow q) \wedge p] \rightarrow q$
T	T	T	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

NOTES:

- We always connect all hypotheses with a \wedge .
- The argument is valid if and only if all the truth values in the last column are true.

An **invalid argument** is an argument in which the conclusion can be false when its hypothesis is true.

$$\begin{array}{l} \text{If it is raining, then the streets are wet.} \\ \text{The streets are wet.} \\ \hline \text{Therefore, it is raining.} \end{array}$$

Let p = it is raining and let q = the streets are wet. Then symbolically, the above argument can be represented as:

$$\frac{p \rightarrow q}{q} \\ p$$

Again, we determine the validity of this argument by considering its corresponding truth table:

p	q	$p \rightarrow q$	$(p \rightarrow q) \wedge q$	$[(p \rightarrow q) \wedge q] \rightarrow p$
T	T	T	T	T
T	F	F	F	T
F	T	T	T	F
F	F	T	F	T

NOTE: The argument is invalid because not all of the truth values in the last column are true.

Examples: Use truth tables to determine if the following arguments are valid or invalid.

1.

$$\frac{q \wedge \sim p}{\sim p} \\ q$$

2.

If you like Tums, then you'll like Roloids.
You don't like Roloids.

You don't like Tums.

3.

$$\begin{array}{l} \sim p \rightarrow q \\ q \rightarrow p \\ \hline q \end{array}$$

4.

$$\frac{p \wedge \sim q}{\sim p \rightarrow q} \\ \sim (p \vee q)$$

5.

$$\frac{\sim p \vee q}{p \leftrightarrow (q \wedge \sim p)} \\ q \wedge p$$