DNA

We will discuss factors relating to the structure, properties, and uses of deoxyribonucleic acid (DNA).

- Purpose
- Structure
- Replication
- Applications
  - DNA Fingerprints
  - Genetics

Size considerations

The following is known about human DNA.

- If stretched out, the DNA in a single cell would be ∼2 meters long.
- DNA is divided between 23 chromosomes
- Our DNA contains ∼20,000-25,000 genes
  - (Based on Human Genome Project - text states ∼100,000)

DNA's Purpose

DNA contains our "genetic code" - inherited physical traits.

- What is known.
  Certain nucleotide sequences carry the "code" for synthesis of proteins
- What is unknown.
  98% DNA is either "junk" or (more likely) we don't what is does.

See article by Tom Bethell in
The American Spectator, March 2004.

DNA Structure

DNA is composed of four types of nucleotides:



For more information, see Nucleic Acids Handout.

Base Pairs



Adenine always pairs with Thymine (A-T pair).
Cytosine always pairs with Guanine (C-G pair).

Double Helix



Replication

Since each nucleotide forms a “match” with only a single type of nucleotide, if one strand is known, the other can be determined.

–A–	–C–	–G–	–T–	–A–	–A–	–T–	–T–
–T–	–G–	–C–	–A–	–T–	–T–	–A–	–A–

Polymerase Chain Reaction

PCR is a method for synthesizing identical copies of DNA from a very small sample.
The General procedure is as follows.

- A DNA molecule is heated to separate into two strands.
- Free nucleotides (and primer) added to synthesize complementary strands.
- Repeat as needed.

DNA Doubling

$$\begin{array}{c} --A-C-G-T-A-A-A-G-G-G-- \\ --T-G-C-A-T-T-T-C-C-C-- \\ \Downarrow \\ --A-C-G-T-A-A-A-G-G-G-- \\ + \\ --T-G-C-A-T-T-T-C-C-C-- \\ \Downarrow \\ --A-C-G-T-A-A-A-G-G-G-- \\ --T-G-C-A-T-T-T-C-C-C-- \\ + \\ --A-C-G-T-A-A-A-G-G-G-- \\ --T-G-C-A-T-T-T-C-C-C-- \end{array}$$

PCR - Exponential Growth

Which would you rather have, a $1,000,000 or a penny a day, doubled every day for a month?

$$1 \longrightarrow 2 \longrightarrow 4 \longrightarrow 16 \longrightarrow 32 \longrightarrow \ldots$$

- After 10 cycles, 1024 times as much.
- After 20 cycles, $\sim$1 million times as much.
- After 30 cycles, $\sim$1 billion times as much.

In PCR, $\sim$1-2 minutes/cycle.

DNA Fingerprinting

- Collect DNA from “crime scene”.
- Use PCR to create large quantities of identical copies of DNA.
- Cut into small pieces using Restriction Enzymes
- Separate pieces based on size using Gel Electrophoresis

DNA Fingerprinting

(From St. Anselm College)

Paternity Testing



(From PBS website)

RNA

RNA is used primarily for protein synthesis.
There are three different types of RNA.

- messenger RNA (mRNA) - transcribes (copies) genetic code from DNA.
- transfer RNA (tRNA) - transports amino acids to the site of protein synthesis.
- ribosomal RNA (rRNA) - responsible (along with some protein) for translating the DNA code into an amino acid sequence for protein synthesis.

Protein Coding

Problem:
* Only four nucleotides to code for 20 amino acids
Solution:
* Use more than one.
(Analogous to 26 letter alphabet ⟶ all possible words)

It requires a three-nucleotide sequence to code for all 20 amino acids.
(There are actually 64 ways to arrange three nucleotides)

Analogy



Insulin Production

Insulin contains 51 amino acids.
Laboratory synthesis too difficult.
Insulin obtained from:

- Slaughterhouses (Cows or Pigs)
- Bacteria using Recombinant DNA techniques (p 494)

Genetic Disease

- Relatively rare
- Error in DNA ⟶ synthesis of “bad” protein
- Usually “triggered” by something
- Examples include Down’s syndrome, ALS, ...

Chromosomes

- Humans contain 23 pairs of chromosomes.
- Children receive half of their chromosomes from each parent.
- Genetics often analyzed using **Punnett Squares**.

Genetics

Example, if both parents are **heterozygous**, they have a dominant (**R**) and recessive (**r**) gene for a particular trait:

		Dad	
		R	**r**
Mom	**R**	RR	Rr
	r	Rr	rr

25% of children will have recessive trait (**rr**).
75% will show characteristic of dominant trait (**R**).

X-linked disease - Mom

Mother 'normal', but has recessive gene (x) for disease. For example, this occurs with hemophilia, which is a disease that prevents blood from clotting.

		Dad	
		X	**Y**
Mom	**X**	XX	XY
	x	Xx	xY

X-linked disease - Dad

Dad carries recessive gene (x) for disease. Because he has only one X chromosome, he also carries disease.

		Dad	
		x	**Y**
Mom	**X**	xX	XY
	X	xX	XY

Y-linked Genetics

Traits on "Y" chromosome

- Comes from father
- Passed to all sons (∼100%)
- Never given to daughters
- Almost always expressed (X ≠ Y)
- Diseases relatively rare

Non-Dominant Traits

For some traits, combinations of different genes gives rise not to the dominant trait but a combination (**incomplete dominance**).

- Ex. – For some flowers, R_{red} + r_{yellow} ⇒ Rr_{orange}

Some genetic traits, such as skin color, are governed by more than one type of gene (**polygenic**).

- Ex. – Skin color in humans determined by at least three different genes, whose effects are additive.

Skin Color

For simplicity, assume only two genes (A & B) determine skin color.

		Dad			
		AB	Ab	aB	ab
Mom	AB		AABb	AaBB	AaBb
	Ab	AABb	AAbb	AaBb	Aabb
	aB	AaBB	AaBb	aaBB	aaBb
	ab	AaBb	Aabb	aaBb	aabb