

**Definition:**

- **Quadratic Equation:** is an equation that can be written in the form

$$ax^2 + bx + c = 0,$$

where  $a$ ,  $b$ , and  $c$  are real numbers,  $a \neq 0$ .

**Steps for solving a quadratic equation by completing the square :**

1. Isolate the constant on one side.
2. Make sure that the coefficient in front of the squared term is a *positive* one. If it is not, divide every term of the equation by this coefficient.
3. Once the coefficient on the squared term is a positive one, take one-half the coefficient of  $x$  and square this result. Namely,  $(\frac{1}{2} \cdot \text{coeff of } x)^2$ .
4. Add the result of Step 3 to both sides.
5. Factor the side containing the variables as a perfect square.
6. Solve the resulting equation using the square root property.

**Important Properties:**

- **Square Root Property:** If  $c$  is a positive number and if  $x^2 = c$ , then

$$x = \sqrt{c} \quad \text{or} \quad x = -\sqrt{c}.$$

(This can be written in one statement as  $\pm\sqrt{c}$ .) In other words, when solving a quadratic equation by the square root property, we want both the positive and negative square roots.

- Completing the square can be used to solve *any* quadratic equation.

**Common Mistakes to Avoid:**

- While completing the square, do not proceed to step three until the coefficient on the squared term is a POSITIVE one.
- Do NOT forget the  $\pm$  when taking square roots.
- If the solution can be simplified, you must do so. For example,  $x = -5 \pm 2$  becomes  $x = -5 + 2 = -3$  and  $x = -5 - 2 = -7$ .

## PROBLEMS

Solve the following equations by completing the square.

1.  $x^2 + 4x = 5$

$$\begin{aligned}
 x^2 + 4x &= 5 \\
 \left(\frac{1}{2} \cdot 4\right)^2 &= (2)^2 = 4 \\
 x^2 + 4x + 4 &= 5 + 4 \\
 (x + 2)^2 &= 9 \\
 \sqrt{(x + 2)^2} &= \sqrt{9} \\
 x + 2 &= \pm 3 \\
 x &= -2 \pm 3 \\
 x &= -2 + 3 = 1 \\
 x &= -2 - 3 = -5
 \end{aligned}$$

$$x = 1, \quad x = -5$$

2.  $x^2 + 4x + 1 = 0$

$$\begin{aligned}
 x^2 + 4x + 1 &= 0 \\
 x^2 + 4x &= -1 \\
 \left(\frac{1}{2} \cdot 4\right)^2 &= (2)^2 = 4 \\
 x^2 + 4x + 4 &= -1 + 4 \\
 (x + 2)^2 &= 3 \\
 \sqrt{(x + 2)^2} &= \sqrt{3} \\
 x + 2 &= \pm\sqrt{3} \\
 x &= -2 \pm \sqrt{3}
 \end{aligned}$$

$$x = -2 + \sqrt{3}, \quad x = -2 - \sqrt{3}$$

3.  $x^2 - 8x - 5 = 0$

$$\begin{aligned}
 x^2 - 8x - 5 &= 0 \\
 x^2 - 8x &= 5 \\
 \left(\frac{1}{2} \cdot -8\right)^2 &= (-4)^2 = 16 \\
 x^2 - 8x + 16 &= 5 + 16 \\
 (x - 4)^2 &= 21 \\
 \sqrt{(x - 4)^2} &= \sqrt{21} \\
 x - 4 &= \pm\sqrt{21} \\
 x &= 4 \pm \sqrt{21}
 \end{aligned}$$

$$x = 4 + \sqrt{21}, \quad x = 4 - \sqrt{21}$$

4.  $2x^2 - 12x + 14 = 0$

$$\begin{aligned}
 2x^2 - 12x + 14 &= 0 \\
 2x^2 - 12x &= -14 \\
 x^2 - 6x &= -7 \\
 \left(\frac{1}{2} \cdot -6\right)^2 &= (-3)^2 = 9 \\
 x^2 - 6x + 9 &= -7 + 9 \\
 (x - 3)^2 &= 2 \\
 \sqrt{(x - 3)^2} &= \sqrt{2} \\
 x - 3 &= \pm\sqrt{2} \\
 x &= 3 \pm \sqrt{2}
 \end{aligned}$$

$$x = 3 + \sqrt{2}, \quad x = 3 - \sqrt{2}$$

5.  $4x^2 + 32x - 3 = 0$

$$4x^2 + 32x - 3 = 0$$

$$4x^2 + 32x = 3$$

$$x^2 + 8x = \frac{3}{4}$$

$$\left(\frac{1}{2} \cdot 8\right)^2 = (4)^2 = 16$$

$$x^2 + 8x + 16 = \frac{3}{4} + 16$$

$$(x + 4)^2 = \frac{69}{4}$$

$$\sqrt{(x + 4)^2} = \sqrt{\frac{69}{4}}$$

$$x + 4 = \pm \frac{\sqrt{69}}{2}$$

$$x = -4 \pm \frac{\sqrt{69}}{2}$$

$$x = -4 + \frac{\sqrt{69}}{2}, \quad x = -4 - \frac{\sqrt{69}}{2}$$

6.  $5x^2 + 5x - 15 = 0$

$$5x^2 + 5x - 15 = 0$$

$$5x^2 + 5x = 15$$

$$x^2 + x = 3$$

$$\left(\frac{1}{2} \cdot 1\right)^2 = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$$

$$x^2 + x + \frac{1}{4} = 3 + \frac{1}{4}$$

$$\left(x + \frac{1}{2}\right)^2 = \frac{13}{4}$$

$$\sqrt{\left(x + \frac{1}{2}\right)^2} = \sqrt{\frac{13}{4}}$$

$$x + \frac{1}{2} = \pm \frac{\sqrt{13}}{2}$$

$$x = -\frac{1}{2} \pm \frac{\sqrt{13}}{2}$$

$$x = -\frac{1}{2} + \frac{\sqrt{13}}{2}, \quad x = -\frac{1}{2} - \frac{\sqrt{13}}{2}$$

7.  $3x^2 + 7x = 4$

$$3x^2 + 7x = 4$$

$$x^2 + \frac{7}{3}x = \frac{4}{3}$$

$$\left(\frac{1}{2} \cdot \frac{7}{3}\right)^2 = \left(\frac{7}{6}\right)^2 = \frac{49}{36}$$

$$x^2 + \frac{7}{3}x + \frac{49}{36} = \frac{4}{3} + \frac{49}{36}$$

$$x^2 + \frac{7}{3}x + \frac{49}{36} = \frac{48}{36} + \frac{49}{36}$$

$$x^2 + \frac{7}{3}x + \frac{49}{36} = \frac{97}{36}$$

$$\left(x + \frac{7}{6}\right)^2 = \frac{97}{36}$$

$$\sqrt{\left(x + \frac{7}{6}\right)^2} = \sqrt{\frac{97}{36}}$$

$$x + \frac{7}{6} = \pm \frac{\sqrt{97}}{6}$$

$$x = -\frac{7}{6} \pm \frac{\sqrt{97}}{6}$$

$x = -\frac{7}{6} + \frac{\sqrt{97}}{6}, \quad x = -\frac{7}{6} - \frac{\sqrt{97}}{6}$
--